

CSci 1302 Example of a proof in predicate logic

The example uses the following predicates on integer numbers:

1. $next(x, y)$ is true if $y = x + 1$, false otherwise.
2. $greater(x, y)$ is true if $x > y$, false otherwise.

Below are two proofs of the same mathematical argument: if every number has the next number and the next number of x is greater than x , then there is no greatest number. Note that we need to add to our list of assumptions the property that if x is greater than y , then y is not greater than x . Without this assumption the proof does not go through.

There are two versions of the proof. The first one uses the theorems of predicate logic $\sim\exists x.p(x) \equiv \forall x.\sim p(x)$ and $\sim\forall x.p(x) \equiv \exists x.\sim p(x)$. The other one is a proof by contradiction.

1.	$\forall x.\exists y.next(x, y)$	<i>Assumption</i>
2.	$\forall x.\forall y.next(x, y) \rightarrow greater(y, x)$	<i>Assumption</i>
3.	$\forall x.\forall y.greater(x, y) \rightarrow \sim greater(y, x)$	<i>Assumption</i>
	$\therefore \sim\exists x.\forall y.greater(x, y)$	
4.	$\exists y.next(x_{\forall}, y(x_{\forall}))$	1, $\forall_E(gen)$, y depends on x_{\forall}
5.	$next(x_{\forall}, y_{\exists}(x_{\forall}))$	4, \exists_E
6.	$next(x_{\forall}, y_{\exists}(x_{\forall})) \rightarrow greater(y_{\exists}(x_{\forall}), x_{\forall})$	2, $\forall_E(unknown)$ twice (x_{\forall} appeared before, so it's unknown, not genuine)
7.	$greater(y_{\exists}(x_{\forall}), x_{\forall})$	5, 6, Modus Ponens
8.	$greater(y_{\exists}(x_{\forall}), x_{\forall}) \rightarrow \sim greater(x_{\forall}, y_{\exists}(x_{\forall}))$	3, $\forall_E(unknown)$ twice
9.	$\sim greater(x_{\forall}, y_{\exists}(x_{\forall}))$	7, 8, Modus Ponens
10.	$\exists y.\sim greater(x_{\forall}, y)$	9, \exists_I
11.	$\forall x.\exists y.\sim greater(x, y)$	10, \forall_I
12.	$\forall x.\sim\forall y.greater(x, y)$	11, Thm. $\sim\forall x.p(x) \equiv \exists x.\sim p(x)$
13.	$\sim\exists x.\forall y.greater(x, y)$	12, Thm. $\sim\exists x.p(x) \equiv \forall x.\sim p(x)$

Note that step 10 introduces \exists quantifier for a variable y_{\exists} which depends on x_{\forall} . This is OK. However, introducing the \forall quantifier would not have been allowed.

The same example can be done using a proof by contradiction: assume that there is the greatest number, and find a contradiction.

1.	$\forall x.\exists y.next(x, y)$	<i>Assumption</i>
2.	$\forall x.\forall y.next(x, y) \rightarrow greater(y, x)$	<i>Assumption</i>
3.	$\forall x.\forall y.greater(x, y) \rightarrow \sim greater(y, x)$	<i>Assumption</i>
<hr/>		
	$\sim \exists x \forall y.greater(x, y)$	
4.	$\exists x.\forall y.greater(x, y)$	<i>Assumption</i> (there is the largest number)
5.	$\forall y.greater(x_{\exists}, y)$	4, \exists_E
6.	$\exists y.next(x_{\exists}, y)$	1, $\forall_E(unknown)$
7.	$next(x_{\exists}, y_{\exists})$	6, \exists_E
8.	$next(x_{\exists}, y_{\exists}) \rightarrow greater(y_{\exists}, x_{\exists})$	7, $\forall_E(unknown)$ twice
9.	$greater(y_{\exists}, x_{\exists})$	7, 8, Modus Ponens
10.	$greater(y_{\exists}, x_{\exists}) \rightarrow \sim greater(x_{\exists}, y_{\exists})$	3, $\forall_E(unknown)$ twice
11.	$\sim greater(x_{\exists}, y_{\exists})$	9, 10, Modus Ponens
12.	$greater(x_{\exists}, y_{\exists})$	5, $\forall_E(unknown)$
13.	$\sim \exists x.\forall y.greater(x, y)$	4 – 12, Contradiction lines 11, 12